


आ.ब. २०७९/८० मा
धानवालीमा देखिएका समस्याहरु

ब्याक्टेरियल पात डडुवा Bacterial leaf Blight

रोग: ब्याक्टेरिया

ब्यवस्थापन

कपर अक्सिक्लोराईड ५०% डब्लु पी १५-२० ग्राम र स्ट्रेप्टोमाईसिन सल्फेट ९०% +
टेट्रासाईक्लिन हाईड्रोक्लोराईड १०% ३ ग्राम १ टंकी पानीमा मिसाएर छर्ने ।



ब्याक्टेरियल बाला डडुवा Bacterial Panicle Blight

रोग: ब्याक्टेरिया *Burkholderia glumae* or *B. gladioli*

यस रोगको उपचारका लागि कुनैपनी विषादी नभए पनि कपर अक्सिक्लोराईड ५०% डब्लु पी १५-२० ग्राम र स्ट्रेप्टोमाईसिन सल्फेट ९०% + टेट्रासाईक्लिन हाईड्रोक्लोराईड १०% एस पि २ ग्राम १ टंकी पानीमा मिसाएर छर्ने ।



Direction for use: Please see the literature.
उपयोग के लिये कृपया निर्देश पत्र देखें।

स्ट्रेप्टोसायक्लिन® STREPTOCYCLINE®

FOR AGRICULTURAL USE ONLY

COMPOSITION :

Streptomycin Sulphate IP 90% w/w
Tetracycline Hydrochloride IP 10% w/w

Net Content 6 g

STORE IN A COOL DRY PLACE AWAY FROM SUNLIGHT.
ठंडी जगह, सुखी जगह पर और सूर्यप्रकाश से दूर रखिये।

KEEP OUT OF THE REACH OF CHILDREN.
बच्चों के हाथों से दूर सुरक्षित रखिये।

Mfg. Lic. No. L01M0035
Reg. No. CIR-592/61 Streptocycline (SP)-2

Batch No.

Mfg. Date

उत्पादन तिथि

Expiry Date

व्यक्ति की तिथि

Maximum Retail Price Rs.

(Inclusive of all taxes)

सर्व अधिकतम मूल्य (सर्व कर सहित)



CAUTION



TATA BLITOX® 50 W COPPER OXYCHLORIDE 50 % WP (containing 50% w/w copper content) CONTACT FUNGICIDE

Chemical Composition : Copper oxychloride 50% w/w, Wetting agent (5% w/w), Dispersing agent (5% w/w), Filler (30% w/w), Total 100% w/w.

Recommendations : It is recommended to control leaf spot & fruit rot of banana, late & early blight of potato & potato leaf spot & late rot of potato, downy mildew of grapes and disease of paddy, cotton, citrus, coffee, tea and tobacco crops.

Precautions : Avoid inhalation & skin contact. Wash contaminated skin & clothes. Do not eat, drink or smoke during application. Destroy empty containers after use.

Antidote : Empty stomach by gastric lavage with milk or preferably w/v sol. of potassium permanganate. Administer egg white and other demulcent. Monitor electrolyte and fluid balance. Morphine or pethidine may be given to control pain. Dose of potassium permanganate 10-40 mg/day (max. 1-2 g/day) divided into four dosages should be administered for five days. Omeprazole can be given 1 mg/kg every four hours for first four days and 1 mg/kg every 12 hours for a total of 10 days. Give water first four hours as

per poisoning to obtain maximum benefit. Omeprazole is administered intravenously. **Direction for use :** Please read enclosed leaflet before use. **Caution :** Not to be used on crops other than specified on this label.

- Keep in a cool, dry place, away from heat or open flame.
- KEEP OUT OF REACH OF CHILDREN. READ THE ENCLOSED LEAFLET CAREFULLY BEFORE USE.



TATA INDIA LIMITED
A TATA Enterprise





धानको पातको फेद डढुवा Sheath Blight

ब्यवस्थापन

भेलिडामाईसिन ३% L @ २-३ मि.लि. प्रति १ लिटरपानीमा मिसाएर छर्ने । वा

propiconazole 10.7% + Tricyclazole 34.2% SE systemic fungicide 2 ml per litre water



धानको खैरो थोप्ले र मरुवा रोग **Brown spot and Blast**

ब्यवस्थापन

Azoxystrobin 25%+ Thifluzamide 20% SC @ २-३ मि.लि. प्रति १ टंकी पानीमा मिसाएर छर्ने ।

propiconazole 10.7% + Tricyclazole 34.2% SE systemic fungicide 2 ml per litre water



धानको पातको फेद डडुवा Sheath Blight र सेतो झिंगा whitefly

भेलिडामार्इसिन ३% L @ २-३ मि.लि. प्रति १ टंकी पानीमा मिसाएर छर्ने ।

propiconazole 10.7% + Tricyclazole 34.2% SE systemic fungicide 2 ml per litre water
सेतो झिंगाको लागि Dimethoate 30% EC वा Imidacloprid 17.8% SL @1.5 ml per litre



